



# Soils Map



Soils data provided by USDA and NRCS.



State: **Indiana**  
 County: **Blackford**  
 Location: **9-23N-11E**  
 Township: **Jackson**  
 Acres: **239.16**  
 Date: **9/16/2026**



**HALDERMAN**  
 REAL ESTATE & FARM MANAGEMENT

Maps Provided By:



© AgriData, Inc. 2025

www.AgriDataInc.com



Area Symbol: IN009, Soil Area Version: 28

| Code             | Soil Description                                                          | Acres | Percent of field | Non-Irr Class Legend | Non-Irr Class *c | Corn Bu | Grass legume hay Tons | Grass legume pasture AUM | Pasture AUM | Soybeans Bu | Winter wheat Bu | *n NCCPI Soybeans |
|------------------|---------------------------------------------------------------------------|-------|------------------|----------------------|------------------|---------|-----------------------|--------------------------|-------------|-------------|-----------------|-------------------|
| GleB2            | Glynwood silt loam, end moraine, 1 to 4 percent slopes, eroded            | 80.24 | 33.7%            |                      | Ile              | 123     | 4                     | 8                        |             | 42          | 55              | 38                |
| BIA              | Blount-Glynwood, thin solum complex, 0 to 3 percent slopes                | 61.30 | 25.6%            |                      | Ilw              | 131     | 5                     |                          | 9           | 44          | 59              | 48                |
| Pm               | Pewamo silty clay, 0 to 2 percent slopes                                  | 48.15 | 20.1%            |                      | Ilw              | 153     | 5                     | 10                       |             | 43          | 62              | 66                |
| GlgB2            | Glynwood silt loam, ground moraine, 1 to 4 percent slopes, eroded         | 25.68 | 10.7%            |                      | Ile              | 123     | 4                     | 8                        |             | 42          | 55              | 38                |
| St               | Saranac silty clay, 0 to 2 percent slopes, frequently flooded             | 11.76 | 4.9%             |                      | Illw             | 128     | 4                     | 8                        |             | 45          | 6               | 71                |
| GlpC3            | Glynwood clay loam, 6 to 12 percent slopes, severely eroded               | 8.22  | 3.4%             |                      | IVe              | 112     | 4                     | 7                        |             | 29          | 49              | 29                |
| GlyC3            | Glynwood-Mississinewa clay loams, 6 to 12 percent slopes, severely eroded | 3.81  | 1.6%             |                      | IVe              | 105     | 4                     | 7                        |             | 27          | 47              | 26                |
| Weighted Average |                                                                           |       |                  |                      | 2.15             | 130.7   | 4.5                   | 6.3                      | 2.3         | 42.2        | 54.7            | *n 47.3           |

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.