

CIVIL CONSTRUCTION DRAWINGS

RESIDENTIAL DRIVEWAY DESIGN AND GRADING PLAN

11002 TWIN SPRUCE ROAD, GOLDEN
COUNTY OF JEFFERSON, STATE OF COLORADO

LEGEND:

- PROPOSED DRIVEWAY (PRIVATE)
- AREA TO BE REVEGETATED
- TOTAL AREA = 21,424 SF

REVEGETATION PLAN:

EROSION CONTROL MEASURES: EROSION CONTROL BLANKET TO BE UTILIZED ON ALL SLOPES GREATER THAN 4:1 IN ADDITION TO SEEDING.

SEED MIX: LOAMY OR CLAYEY SOILS MIX (SEE SHEET C4 FOR MIXTURE DETAILS, SEEDING RATES AND METHOD OF SEEDING)

SEEDBED PREP: TO THE MAXIMUM EXTENT FEASIBLE, TOPSOIL THAT IS REMOVED DURING CONSTRUCTION ACTIVITY SHALL BE CONSERVED FOR LATER USE ON AREAS REQUIRING REVEGETATION AND LANDSCAPING. REFER TO MILE HIGH FLOOD DISTRICT NONSTRUCTURAL BMP EC-2 FOR MORE DETAILS ON SEED BED PREPARATION.

SEEDING DATES: PERMANENT SEEDING SHALL BE PERFORMED PROMPTLY AFTER REACHING FINAL GRADE (TYPICALLY WITHIN 14 DAYS). REFER TO MILE HIGH FLOOD DISTRICT NONSTRUCTURAL BMP EC-2 FOR MORE DETAILS ON SEEDING DATES WITH THE HIGHEST PROBABILITY OF PROVIDING SUFFICIENT VEGETATIVE COVER

MULCHING RATES AND APPLICATION: SEE MULCHING DETAIL ON SHEET C4 FOR DETAILS ON RATES AND METHODS OF APPLICATION.

Temporary and Permanent Seeding (TS/PS) EC-2

Seeding rates for the highest percent probability of germination are generally in the spring from April through early May and in the fall after the first of September and the ground freezes. If the area is irrigated, seeding may occur in summer months, as well. See Table TS/PS-5 for appropriate seeding rates.

Table TS/PS-1: Minimum Drift Seeding Rates for Various Temporary Annual Grasses

Species (Common name)	Growth Season	Pounds of Pure Live Seed (PLS) per Acre	Planting Depth (inches)
1. Oat	Cool	25-35	1-2
2. Spring wheat	Cool	25-35	1-2
3. Spring barley	Cool	25-35	1-2
4. Annual ryegrass	Cool	10-15	1/2
5. Millet	Warm	3-10	1/2-1
6. Sorghum	Warm	5-10	1/2-1
7. Sorghum	Warm	5-10	1/2-1
8. Winter wheat	Cool	20-35	1-2
9. Winter barley	Cool	20-35	1-2
10. Winter rye	Cool	20-35	1-2
11. Trifolium	Cool	20-40	1-2

Hydraulic seeding may be substituted for drilling only where slopes are steeper than 1:1 or where erosion potential is low. When hydraulic seeding is used, hydraulic mulching should be applied as a separate operation, when practical, to prevent the seeds from being incorporated in the wash.

* See Table TS/PS-5 for seeding rates. Irrigation, if consistently applied, may extend the use of cool season species during the summer months.

* Seeding rates should be doubled if used as broadcast, or increased by 50 percent if done using a blower drill or by hydraulic seeding.

Temporary and Permanent Seeding (TS/PS) EC-2

Table TS/PS-2: Minimum Drift Seeding Rates for Permanent Grasses (cont.)

Species Name	Establishment Season	Seeding Rate (lb/acre)	Planting Depth (inches)	Perennial PLS/acre
Blue grass	Summer/autumn	100,000	1/2	0.5
Compendium	Summer/autumn	100,000	1/2	1.0
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5

* All of the above seeding rates are based on soil testing followed by appropriate seed mix. These rates should be doubled if used as broadcast, or increased by 50 percent if done using a blower drill or by hydraulic seeding. See Table TS/PS-5 for seeding rates.

* If ERM is to be irrigated, the maximum seed rates should be doubled.

* Cool season species should be used on slopes steeper than 1:1.

* Use a minimum of 1/2 inch of the above rates for 1:1 to 2:1 slopes of 1/2 inch of the above rates for 2:1 to 3:1 slopes.

EC-2 Temporary and Permanent Seeding (TS/PS)

Table TS/PS-2: Minimum Drift Seeding Rates for Permanent Grasses

Species Name	Establishment Season	Seeding Rate (lb/acre)	Planting Depth (inches)	Perennial PLS/acre
Blue grass	Summer/autumn	100,000	1/2	0.5
Compendium	Summer/autumn	100,000	1/2	1.0
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5
Hard fescue	Summer/autumn	100,000	1/2	0.5

* All of the above seeding rates are based on soil testing followed by appropriate seed mix. These rates should be doubled if used as broadcast, or increased by 50 percent if done using a blower drill or by hydraulic seeding. See Table TS/PS-5 for seeding rates.

* If ERM is to be irrigated, the maximum seed rates should be doubled.

* Cool season species should be used on slopes steeper than 1:1.

* Use a minimum of 1/2 inch of the above rates for 1:1 to 2:1 slopes of 1/2 inch of the above rates for 2:1 to 3:1 slopes.

EC-2 Temporary and Permanent Seeding (TS/PS)


Table TS/PS-3: Seeding Dates for Annual and Perennial Grasses

Seeding Dates	Annual Grasses (lb/acre)	Perennial Grasses (lb/acre)
January - March 15	4	1.25
March 15 - April 30	4	1.25
April 30 - May 15	4	1.25
May 15 - June 30	4	1.25
June 30 - July 15	4	1.25
July 15 - August 31	4	1.25
August 31 - September 30	4	1.25
September 30 - October 31	4	1.25
October 31 - November 31	4	1.25
November 31 - December 31	4	1.25


Mulch
Cover seeded areas with mulch or an appropriate erosion control product to prevent establishment of vegetation. Mulch must be applied, setting or use of a mulch machine. See the Mulching BMP Fact Sheet for additional guidance.

Maintenance and Removal
Monitor and observe seeded areas to identify areas of poor growth or areas that fail to germinate. Remove and reseed these areas, as needed.
An area that has been permanently seeded should have a good stand of vegetation within one growing season. If removal and within three growing seasons without irrigation in Colorado. Harvest portions of the site that fail to germinate or contain bare after the first growing season.
Seeded areas may require irrigation, particularly during extended dry periods. Targeted weed control may also be necessary.
Protect seeded areas from construction equipment and vehicle access.

TEMPORARY AND PERMANENT SEEDING DETAILS



VICINITY MAP
1" = 2000'



REVEGETATION PLAN



BASELINE

Engineering • Planning • Surveying
102 N. RIBEY DRIVE, SUITE 210 • GOLDEN, COLORADO 80403
P: 303.940.9966 • F: 303.940.9965 • www.baselinecorp.com

DESIGNED BY: NN
DRAWN BY: LR
CHECKED BY: NN

DATE: 2/12/23

PREPARED BY: NN

PER COUNTY SUFFICIENCY REVIEW

REVISION DESCRIPTION

JEFFERSON COUNTY

11002 TWIN SPRUCE ROAD

UNINCORPORATED

REVEGETATION PLAN

DARIUSZ STACHOWIAK

GOLDEN, COLORADO

PREPARED UNDER THE DIRECT SUPERVISION OF

FOR AND ON BEHALF OF

INITIAL SUBMITTAL: XX/XX/XX

DRAWING SIZE: 24" X 36"

SURVEY FIRM: SURVEY DATE: XX/XX/XX

JOB NO.: C03547

DRAWING NAME: C03547-REVEGETATION.dwg

SHEET 8 OF 10

C8