



Soils Map



Soils data provided by USDA and NRCS.



State: **Indiana**
County: **Benton**
Location: **22-24N-7W**
Township: **Bolivar**
Acres: **283.67**
Date: **2/12/2026**



Maps Provided By:



© AgriData, Inc. 2025

HALDERMAN
REAL ESTATE & FARM MANAGEMENT



www.AgriDataInc.com

Area Symbol: IN007, Soil Area Version: 28									
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Non-Irr Class *c	Corn Bu	Soybeans Bu	*n NCCPI Overall	*n NCCPI Corn
Dp	Darroch silt loam, till substratum	98.46	34.7%		IIw	160	49	80	80
Dx	Drummer silty clay loam, stratified sandy substratum	52.36	18.5%		IIw	175	49	91	91
Ch	Chalmers silty clay loam	42.74	15.1%		IIw	190	54	81	81
BaB2	Barce loam, 2 to 6 percent slopes, eroded	19.37	6.8%		Ile	141	46	76	76
FpB2	Foresman silt loam, till substratum, 1 to 5 percent slopes, eroded	13.62	4.8%		Ile	144	47	75	75
Do	Darroch silt loam	11.77	4.1%		IIw	158	49	86	83
Cu	Crane loam, till substratum	11.01	3.9%		IIw	150	46	79	79
GIA	Gilboa silt loam, 0 to 2 percent slopes	8.69	3.1%		IIw	155	48	88	88
Wb	Warners variant silty clay, undrained	6.64	2.3%		Vw			33	33
GIB	Gilboa silt loam, 2 to 4 percent slopes	6.06	2.1%		Ile	152	47	87	87
Sh	Selma silty clay loam, till substratum	5.93	2.1%		IIw	175	49	84	84
BbA	Barce silt loam, 0 to 2 percent slopes	4.72	1.7%		I	148	49	89	89
BdB2	Billett sandy loam, 2 to 6 percent slopes, eroded	1.68	0.6%		IIIe	122	41	67	67
OIA	Odell silt loam, 0 to 2 percent slopes	0.40	0.1%		IIw	153	47	76	76
Cm	Comfrey silty clay loam, sandy substratum, frequently flooded	0.22	0.1%		Vw	145	37	50	50
Weighted Average					2.06	160.6	48.1	*n 81.3	*n 81.2

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.